



Dr. Stansifer – Pediatric Ear Tubes – (402) 397-0670

- Your child may be fussy on the day of surgery from anesthesia
- There may be some ear discharge (clear, yellow and/or blood tinged) for a few days
- A low grade fever of 100-101° is common for several days after surgery
You can treat with Tylenol (acetaminophen) and/or Motrin (ibuprofen) as needed
- **Ciprofloxacin-dexamethasone ear drops**
 - Start today: 4 drops to both ears, 3 times daily for ____ days
 - Shake and warm bottle in your hand before use
 - Massage tab of cartilage in front of ear after applying drops
- Precautions
 - There are no activity, exercise, or flying restrictions
 - No ear plugs are needed for baths, showers, swimming pools
 - Ear plugs (kids swimming reusable silicone ear plugs) & headwrap (i.e. Ear Band-It) **are needed** for ocean, river, lake water
- Use the ear drops for 4-7 days for future episodes of ear drainage
 - Please call the office if the drainage does not resolve with the drops



Dr. Stansifer – Pediatric Ear Tubes – (402) 397-0670

- Your child may be fussy on the day of surgery from anesthesia
- There may be some ear discharge (clear, yellow and/or blood tinged) for a few days
- A low grade fever of 100-101° is common for several days after surgery
You can treat with Tylenol (acetaminophen) and/or Motrin (ibuprofen) as needed
- **Ciprofloxacin-dexamethasone ear drops**
 - Start today: 4 drops to both ears, 3 times daily for ____ days
 - Shake and warm bottle in your hand before use
 - Massage tab of cartilage in front of ear after applying drops
- Precautions
 - There are no activity, exercise, or flying restrictions
 - No ear plugs are needed for baths, showers, swimming pools
 - Ear plugs (kids swimming reusable silicone ear plugs) & headwrap (i.e. Ear Band-It) **are needed** for ocean, river, lake water
- Use the ear drops for 4-7 days for future episodes of ear drainage
 - Please call the office if the drainage does not resolve with the drops